

## SHORE CARDIOLOGY CONSULTANTS

---

ALI R. MOOSVI, MD, FACP, FACC  
TODD S. COHEN, DO, FACC  
MICHAEL G. DEVITA, DO, FACC

1640 HWY. 88, SUITE 201  
BRICK, NEW JERSEY 08724  
TELEPHONE (732) 840-1900  
FAX (732) 840-0355  
[www.shorecardiology.com](http://www.shorecardiology.com)

### **Cardiac Catheterization Pre-Procedure Instructions**

You are required to have someone drive you to the facility. You may not drink any alcohol, drive a car, or operate any heavy machinery for the next 24 hours.

#### **What to bring with you:**

- Insurance Cards
- A list of your medications (including dosage) or medications in their bottles

#### **Medication Instructions:**

- Stop taking Coumadin/Warfarin four days prior to procedure
- Stop Pradaxa 24 hours prior to procedure
- Do not take Glucophage/Glucoavance the day of the procedure. Do not take Metformin the day of your procedure and two days after the procedure.
- Take ½ the dose of insulin on the day of your procedure
- Take all other morning medications

#### **Eating/Drinking:**


- Do not eat/drink anything after midnight of your procedure
- Take water with morning medications on the day of the procedure

#### **Day of procedure instructions:**

- If you wear contact lenses, please leave them at home and wear glasses
- Do not bring any valuables

#### **What to expect:**

- Potential delays due to unscheduled emergencies
- A potential overnight stay based on the outcome of your procedure



## SHORE CARDIOLOGY CONSULTANTS

---

ALI R. MOOSVI, MD, FACP, FACC  
TODD S. COHEN, DO, FACC  
MICHAEL G. DEVITA, DO, FACC

1640 HWY. 88, SUITE 201  
BRICK, NEW JERSEY 08724  
TELEPHONE (732) 840-1900  
FAX (732) 840-0355  
www.shorecardiology.com

### **Cardiac Catheterization Discharge Instructions**

You are required to have an adult with you for the next 24 hours. You may not drink any alcohol, drive a car, or operate any heavy machinery for the next 24 hours.

**Call if you have any of the following:** Difficulty breathing or swallowing, chest pain or return of heart symptoms; prolonged vomiting lasting greater than 12 hours and not allowing you to resume your medications and / or regular diet; severe headache unrelieved by Tylenol; redness or swelling at the IV site; or any questions or concerns.

**Additional Instructions:** If you note any signs of bleeding (soaking through the bandage, bulging under the skin, or localized increasing pain), apply direct pressure to the area, call your doctor AND go to the emergency room for evaluation. Anyone who undergoes a procedure and receives a drug eluting stent MUST stay on Plavix or Effient for a minimum of one year, aspirin daily for a lifetime. Do NOT stop these medications for any reason unless cleared with our office directly.

**Activity:** After leaving the hospital, you should rest as much as possible for 24 hours, standing and walking only as necessary (such as going to the bathroom). Avoid heavy lifting and exercise for 7 – 10 days, and minimize stair climbing.

**Medications:** Take as directed by your physician.

**Diet:** Resume your previous diet, except where specified otherwise by your doctor. Drink extra fluids for the next 12 hours (up to 2-3 quarts of water or juice). Please do not drink alcoholic beverages for 24 hours after your procedure.

**Wound Care / Hygiene:** You may shower as soon as you wish. Do not rub the catheterization site in the groin or arm, gently wash and pat dry.

**Sexual Activity:** Refrain from sexual intercourse for the next 24 hours.

**Follow Up in the office:** With Nurse Practitioner or call (732) 840-1900 for an appointment to be seen in 2 weeks.